

General ICU

Name	Telephone	Who is your recruiter?	Date

Instructions: This checklist is meant to serve as a general guideline for our client facilities as to the level of your skills within your nursing specialty. Please use the scale below to describe your experience/expertise in each area listed below.

Proficiency :- 1 = Never Performed
Frequency :- 1 = Never

2 = Limited Experience
2 = Sometimes

3 = Comfortable Performing
3 = Often

4 = Proficient
4 = Always

General ICU Pharmacology

Medications:

	Proficiency				Frequency			
Propofol (Diprivan)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Valium (Diazepam)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Verapamil (Calan, Isoptin, Verelan)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Vecuronium	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Ventolin (Albuterol)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Versed	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Management of Epidural Administration	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Alupent (Metaproterenol)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Amicar	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Aminophylline (Theophylline)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Amiodarone (Cordarone)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Atropine	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
AquaMephyton (Vitamin K)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Apresoline	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Ativan	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Bretylium (Bretylol)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Corticosteroids	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Decadron (Dexamethasone)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Digoxin (Lanoxin)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Dilantin (Phenytoin)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Diltiazem (Cardizem)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Dobutamine (Dobutrex)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Dopamine (Intropin)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Epinephrine (Adrenalin)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Esmolol (Brevibloc)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Fentanyl	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Furosemide (Lasix)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4

Heparin Drip	☐ 1	☐ 2	☐ 3	☐ 4
Inderal (Propranolol)	☐ 1	☐ 2	☐ 3	☐ 4
Inocor (Amrinone)	☐ 1	☐ 2	☐ 3	☐ 4
Kayexalate	☐ 1	☐ 2	☐ 3	☐ 4
Ketamine	☐ 1	☐ 2	☐ 3	☐ 4
Lactulose (Cephulac)	☐ 1	☐ 2	☐ 3	☐ 4
Levophed	☐ 1	☐ 2	☐ 3	☐ 4
Lidocaine	☐ 1	☐ 2	☐ 3	☐ 4
Metoprolol (Lopressor)	☐ 1	☐ 2	☐ 3	☐ 4
Neosynephrine	☐ 1	☐ 2	☐ 3	☐ 4
Nimbex	☐ 1	☐ 2	☐ 3	☐ 4
Nipride (Nitroprusside)	☐ 1	☐ 2	☐ 3	☐ 4
Nitroglycerin (Tridil)	☐ 1	☐ 2	☐ 3	☐ 4
Norepinephrine	☐ 1	☐ 2	☐ 3	☐ 4
Pentobarbital	☐ 1	☐ 2	☐ 3	☐ 4
Pitressin (Vasopressin)	☐ 1	☐ 2	☐ 3	☐ 4
Precedex	☐ 1	☐ 2	☐ 3	☐ 4
Procainamide (Pronestyl)	☐ 1	☐ 2	☐ 3	☐ 4
Sodium Bicarbonate	☐ 1	☐ 2	☐ 3	☐ 4
Succinylcholine	☐ 1	☐ 2	☐ 3	☐ 4

☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4

Provision of Care

Proficiency

Frequency

Perform calculations for weight-based dosages	☐ 1	☐ 2	☐ 3	☐ 4
Perform calculations for oral & parenteral dosages	☐ 1	☐ 2	☐ 3	☐ 4
Assess for drug toxicity	☐ 1	☐ 2	☐ 3	☐ 4
Administer reversal agents when indicated (Narcan, Romazicon, etc.)	☐ 1	☐ 2	☐ 3	☐ 4
Administer/provide safe replacement (isotonic, hypotonic, hypertonic, crystalloid)	☐ 1	☐ 2	☐ 3	☐ 4
Titrate vasopressors to effect/parameters	☐ 1	☐ 2	☐ 3	☐ 4
Assess for therapeutic response	☐ 1	☐ 2	☐ 3	☐ 4
Perform & manage dosage drip calculations	☐ 1	☐ 2	☐ 3	☐ 4
Barbiturate induced coma	☐ 1	☐ 2	☐ 3	☐ 4

☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4

Endocrine

Provision of Care for:

Proficiency

Frequency

SIADH	☐ 1	☐ 2	☐ 3	☐ 4
Addison's Disease	☐ 1	☐ 2	☐ 3	☐ 4
Diabetes mellitus	☐ 1	☐ 2	☐ 3	☐ 4
Diabetic ketoacidosis	☐ 1	☐ 2	☐ 3	☐ 4

☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4

Diabetes Insipidus	☐ 1	☐ 2	☐ 3	☐ 4
Hyperthyroidism	☐ 1	☐ 2	☐ 3	☐ 4
Hypothyroidism	☐ 1	☐ 2	☐ 3	☐ 4
Insulin Shock	☐ 1	☐ 2	☐ 3	☐ 4
Thyroidectomy	☐ 1	☐ 2	☐ 3	☐ 4
Cushing's Syndrome	☐ 1	☐ 2	☐ 3	☐ 4

 1	 2	 3	 4
 1	 2	 3	 4
 1	 2	 3	 4
 1	 2	 3	 4
 1	 2	 3	 4
 1	 2	 3	 4

Procedures:	Proficiency			
Administration of D50	☐ 1	☐ 2	☐ 3	☐ 4
Manage insulin drip	☐ 1	☐ 2	☐ 3	☐ 4
Fingerstick Blood Glucose	☐ 1	☐ 2	☐ 3	☐ 4
Implementation of Sliding Scale	☐ 1	☐ 2	☐ 3	☐ 4

Frequency

☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4

Equipment:	Proficiency			
Blood glucose monitoring devices	☐ 1	☐ 2	☐ 3	☐ 4
Use of Insulin Pen	☐ 1	☐ 2	☐ 3	☐ 4
Use of Lancet	☐ 1	☐ 2	☐ 3	☐ 4

Frequency			
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4

Multisystem				
Provision of Care for:	Proficiency			
Restraints (type, duration, assessment, protocols, and standards)	☐ 1	☐ 2	☐ 3	☐ 4
Therapeutic patient positioning	☐ 1	☐ 2	☐ 3	☐ 4
Implementation of Core Measures	☐ 1	☐ 2	☐ 3	☐ 4
Exploratory laparotomy	☐ 1	☐ 2	☐ 3	☐ 4
Splenectomy	☐ 1	☐ 2	☐ 3	☐ 4
Hypovolemic Shock	☐ 1	☐ 2	☐ 3	☐ 4
Cardiogenic Shock	☐ 1	☐ 2	☐ 3	☐ 4
Septic Shock	☐ 1	☐ 2	☐ 3	☐ 4
Anaphylactic shock	☐ 1	☐ 2	☐ 3	☐ 4
Disseminated Intravascular Coagulation (DIC)	☐ 1	☐ 2	☐ 3	☐ 4
Organ/tissue donation	☐ 1	☐ 2	☐ 3	☐ 4
Sickle Cell Anemia	☐ 1	☐ 2	☐ 3	☐ 4
Multi-system organ failure	☐ 1	☐ 2	☐ 3	☐ 4
Alcohol Overdose	☐ 1	☐ 2	☐ 3	☐ 4
Gunshot Wounds	☐ 1	☐ 2	☐ 3	☐ 4
Cancer	☐ 1	☐ 2	☐ 3	☐ 4
AIDS	☐ 1	☐ 2	☐ 3	☐ 4
Drug Overdose	☐ 1	☐ 2	☐ 3	☐ 4

[illegible]

Emergency Situations:

Follow current ACLS Guidelines

Obtain & utilize crash cart

Administer medications used in emergency situations

Operate defibrillator

Assist with therapeutic hypothermia

Assessment:

Perform comprehensive multi-system assessment

Pain Management:

Assessment of pain level

Assessment of therapeutic response

Care of the patient with:

PICC Line

Epidural anesthesia/analgesia

IV conscious sedation

Patient controlled analgesia

Broviac Catheter

Groshong

Hickman

Portacath

Quinton

Peripheral line/dressing

Labs

Interpretation of:

Blood counts

ABG's

BUN/Creatinine

Coagulation Studies

Electrolytes

Cardiac Enzymes

Blood glucose

Thyroid Panel

Pulmonary**Assessment :**

Interpretation of Oximetry Values

☐ 1 ☐ 2 ☐ 3 ☐ 4

Auscultate breath sounds

☐ 1 ☐ 2 ☐ 3 ☐ 4

Identify adventitious lung sounds

☐ 1 ☐ 2 ☐ 3 ☐ 4

Auscultate for confirmation of ET Tube placement

☐ 1 ☐ 2 ☐ 3 ☐ 4

Implement Ventilator Acquired Pneumonia protocols

☐ 1 ☐ 2 ☐ 3 ☐ 4

Assess and confirm proper Vent settings

☐ 1 ☐ 2 ☐ 3 ☐ 4

Manage patient on mechanical ventilation

☐ 1 ☐ 2 ☐ 3 ☐ 4

Assist and manage patient weaning from artificial pulmonary support

☐ 1 ☐ 2 ☐ 3 ☐ 4

Assist with post-extubation care

☐ 1 ☐ 2 ☐ 3 ☐ 4

Maintain pulmonary stability post-extubation

☐ 1 ☐ 2 ☐ 3 ☐ 4

Application of BIPAP

☐ 1 ☐ 2 ☐ 3 ☐ 4

Application of CPAP

☐ 1 ☐ 2 ☐ 3 ☐ 4

Application of Oximetry

☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4**Provision of Care for:**

Tuberculosis

☐ 1 ☐ 2 ☐ 3 ☐ 4

Acute Pneumonia

☐ 1 ☐ 2 ☐ 3 ☐ 4

ARDS

☐ 1 ☐ 2 ☐ 3 ☐ 4

Chest Trauma

☐ 1 ☐ 2 ☐ 3 ☐ 4

COPD

☐ 1 ☐ 2 ☐ 3 ☐ 4

Cor Pulmonale

☐ 1 ☐ 2 ☐ 3 ☐ 4

Fresh Tracheostomy

☐ 1 ☐ 2 ☐ 3 ☐ 4

Lobectomy

☐ 1 ☐ 2 ☐ 3 ☐ 4

Near Drowning

☐ 1 ☐ 2 ☐ 3 ☐ 4

Pneumonectomy

☐ 1 ☐ 2 ☐ 3 ☐ 4

Pulmonary Edema

☐ 1 ☐ 2 ☐ 3 ☐ 4

Pulmonary Hypertension

☐ 1 ☐ 2 ☐ 3 ☐ 4

Pulmonary Embolism

☐ 1 ☐ 2 ☐ 3 ☐ 4

Respiratory Arrest

☐ 1 ☐ 2 ☐ 3 ☐ 4

Status Asthmaticus

☐ 1 ☐ 2 ☐ 3 ☐ 4

Thoracotomy

☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4☐ 1 ☐ 2 ☐ 3 ☐ 4

Chest Tubes:

	Proficiency				Frequency			
Perform Chest Tube Set-up	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Monitor & record chest tube output	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Identify and troubleshoot clotted chest tube	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Identify and troubleshoot patient bleeding from chest tube	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Identify and troubleshoot presence of air leak	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4

Procedures:

	Proficiency				Frequency			
Perform ET Tube suctioning	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Perform NT (Nasotracheal) suctioning	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Perform suctioning of Tracheostomy	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Collection of Sputum via NT Suction	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Collection of Sputum Specimen via ET (closed system)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Assist with extubation of patient	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4

Neurology:**Assessment :**

	Proficiency				Frequency			
NIH Stroke Scale	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
SS indicating type of stroke (hemorrhagic vs. ischemic)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Cranial Nerves	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Glasgow Coma Scale	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Pathologic Reflexes	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Reflex/Motor Deficits	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Visual/Communication Deficits	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4

Provision of Care for:

	Proficiency				Frequency			
CVA	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
TIA	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Encephalopathy	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Aneurysm Precautions	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Basilar Skull Fracture	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Closed Head Injury	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Coma	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Seizure Disorder	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
DT's	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Encephalitis	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Externalized VP shunts	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4

Procedures:	Proficiency				Frequency			
Obtain Cardiac Output	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Obtain MAP	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
PCWP Measurement	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Obtain CVP Measurement	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Removal of radial arterial line	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Removal of femoral sheath	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Perform Venipuncture	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Obtain lab specimens from A-line	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Obtain lab specimens from Central Line	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Perform 12-lead EKG	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Proper Cardiac Monitor lead placement	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4

Provision of Care for:	Proficiency				Frequency			
AAA (Abdominal Aortic Aneurysm)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Aortic Stenosis	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Cardiomyopathy	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Cardiac Arrest	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Cardiac Catheterization	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Carotid Endarterectomy	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Cardiac Tamponade	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
CHF	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
MI	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Mitral Valve Stenosis	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Myocardial Contusion	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Post cardiac surgery complications	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Pericardial Window	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Post AICD Insertion	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Pericardial Effusion	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Pericarditis	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Fresh Post-op CABG	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
LVAD	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
RVAD	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Fem-Prop Bypass	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4

Assist with:	Proficiency				Frequency			
Set-Up for Cardioversion	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Cardioversion	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Central Line Insertion	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
CAVH-D	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Arterial Line Insertion	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Pulmonary Artery Catheter Insertion	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Chest Tube Insertion	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Chest Tube Removal	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Emergency Tracheostomy	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Pericardiocentesis	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Thoracentesis	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Transesophageal Echocardiogram	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Emergency Intubation	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4

Management of:	Proficiency				Frequency			
Blood and blood product administration	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Blood transfusion reactions	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4

Hemodynamic Monitoring:	Proficiency				Frequency			
Management of Arterial Line	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Management of Pulmonary artery catheter	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Management of CVP Line	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Zero and calibration of transducers	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Assess adequate perfusion with invasive lines	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4

Management of IABP:	Proficiency				Frequency			
Waveform interpretation	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Determination of appropriate timing	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Perfusion assessment	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Recognize IABP complications	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4

Pacemaker:	Proficiency				Frequency			
PPM	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Transvenous	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Transcutaneous	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4

Management of Temporary Pacemaker:**Proficiency****Frequency**

Recognition of failure to sense, capture	☐ 1	☐ 2	☐ 3	☐ 4
Manipulation of TPM settings	☐ 1	☐ 2	☐ 3	☐ 4
Troubleshoot TPM settings	☐ 1	☐ 2	☐ 3	☐ 4

☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4

Identification and interpretation of:**Proficiency****Frequency**

Arterial waveform	☐ 1	☐ 2	☐ 3	☐ 4
Pulmonary artery catheter waveform	☐ 1	☐ 2	☐ 3	☐ 4
PCWP waveform	☐ 1	☐ 2	☐ 3	☐ 4
CVP waveform	☐ 1	☐ 2	☐ 3	☐ 4
SVO2 Data	☐ 1	☐ 2	☐ 3	☐ 4

☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4

Perform calculations for:**Proficiency****Frequency**

PVR	☐ 1	☐ 2	☐ 3	☐ 4
SVR	☐ 1	☐ 2	☐ 3	☐ 4

☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4

Renal:**Assessment :****Proficiency****Frequency**

A-V Fistula/Shunt	☐ 1	☐ 2	☐ 3	☐ 4
Fluid Status	☐ 1	☐ 2	☐ 3	☐ 4

☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4

Provision of Care for:**Proficiency****Frequency**

Acute Renal Failure	☐ 1	☐ 2	☐ 3	☐ 4
CAVH dialysis	☐ 1	☐ 2	☐ 3	☐ 4
Hemodialysis	☐ 1	☐ 2	☐ 3	☐ 4
CRRT	☐ 1	☐ 2	☐ 3	☐ 4
Peritoneal dialysis	☐ 1	☐ 2	☐ 3	☐ 4
Nephrectomy	☐ 1	☐ 2	☐ 3	☐ 4
Renal Transplant	☐ 1	☐ 2	☐ 3	☐ 4
Renal Rejection Syndrome	☐ 1	☐ 2	☐ 3	☐ 4
TURP	☐ 1	☐ 2	☐ 3	☐ 4
Urinary diversion (i.e., ileal, conduit nephrostomy)	☐ 1	☐ 2	☐ 3	☐ 4
Urinary tract infection	☐ 1	☐ 2	☐ 3	☐ 4

☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4

Procedures:**Proficiency****Frequency**

CRRT	☐ 1	☐ 2	☐ 3	☐ 4
CAVH dialysis	☐ 1	☐ 2	☐ 3	☐ 4
Hemodialysis	☐ 1	☐ 2	☐ 3	☐ 4
Bladder Irrigation	☐ 1	☐ 2	☐ 3	☐ 4

☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4

Insertion of Male Foley Catheter	☐ 1	☐ 2	☐ 3	☐ 4
Insertion of Female Foley Catheter	☐ 1	☐ 2	☐ 3	☐ 4
Suprapubic Catheter	☐ 1	☐ 2	☐ 3	☐ 4

☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4
☐ 1	☐ 2	☐ 3	☐ 4

Integumentary:

Procedures:

	Proficiency			
Sterile dressing changes	☐ 1	☐ 2	☐ 3	☐ 4
Wound care/irrigations	☐ 1	☐ 2	☐ 3	☐ 4

	Frequency			
☐ 1	☐ 2	☐ 3	☐ 4	
☐ 1	☐ 2	☐ 3	☐ 4	

Provision of Care for:

	Proficiency			
Burns	☐ 1	☐ 2	☐ 3	☐ 4
Pressure Sores	☐ 1	☐ 2	☐ 3	☐ 4
Staged Decubitus Ulcers	☐ 1	☐ 2	☐ 3	☐ 4
Surgical Wounds w/drains	☐ 1	☐ 2	☐ 3	☐ 4
Traumatic Wounds	☐ 1	☐ 2	☐ 3	☐ 4

	Frequency			
☐ 1	☐ 2	☐ 3	☐ 4	
☐ 1	☐ 2	☐ 3	☐ 4	
☐ 1	☐ 2	☐ 3	☐ 4	
☐ 1	☐ 2	☐ 3	☐ 4	
☐ 1	☐ 2	☐ 3	☐ 4	

Assess for:

	Proficiency			
Skin intact	☐ 1	☐ 2	☐ 3	☐ 4
Stasis Ulcers	☐ 1	☐ 2	☐ 3	☐ 4
Surgical Wound Healing	☐ 1	☐ 2	☐ 3	☐ 4

	Frequency			
☐ 1	☐ 2	☐ 3	☐ 4	
☐ 1	☐ 2	☐ 3	☐ 4	
☐ 1	☐ 2	☐ 3	☐ 4	

Equipment:

	Proficiency			
Air fluidized, low airloss beds	☐ 1	☐ 2	☐ 3	☐ 4

	Frequency			
☐ 1	☐ 2	☐ 3	☐ 4	

Patient Education:

General:

	Proficiency			
Turn, cough, and deep breathing technique	☐ 1	☐ 2	☐ 3	☐ 4
Splinting technique post-op	☐ 1	☐ 2	☐ 3	☐ 4
Use of incentive spirometer	☐ 1	☐ 2	☐ 3	☐ 4
Importance of increasing mobility post-op when applicable	☐ 1	☐ 2	☐ 3	☐ 4

	Frequency			
☐ 1	☐ 2	☐ 3	☐ 4	
☐ 1	☐ 2	☐ 3	☐ 4	
☐ 1	☐ 2	☐ 3	☐ 4	
☐ 1	☐ 2	☐ 3	☐ 4	

Patient Advocacy:

General:

	Proficiency			
Provide care for end-of-life decisions that families face in ICU such as withdrawal of life support	☐ 1	☐ 2	☐ 3	☐ 4
Adhere to legal procedures regarding notification of organ donation provider	☐ 1	☐ 2	☐ 3	☐ 4
Demonstrate Spiritual and Cultural Considerations for the Critical Care patient	☐ 1	☐ 2	☐ 3	☐ 4

	Frequency			
☐ 1	☐ 2	☐ 3	☐ 4	
☐ 1	☐ 2	☐ 3	☐ 4	
☐ 1	☐ 2	☐ 3	☐ 4	

$\bigcirc_1$ $\bigcirc_2$ $\bigcirc_3$ $\bigcirc_4$

O_1 O_2 O_3 O_4

$\bigcirc_1$ $\bigcirc_2$ $\bigcirc_3$ $\bigcirc_4$

$$O_1 \quad O_2 \quad O_3 \quad O_4$$
 $\bigcirc_1 \quad \bigcirc_2 \quad \bigcirc_3 \quad \bigcirc_4$

PPN (peripheral parenteral nutrition)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Removal of N/G Tube	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
TPN and lipids administration	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Utilization of feeding pump	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4

Age Specific

Age Specific:	Proficiency				Frequency			
Infant (Birth to 1 year)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Toddler (1–3 years)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Pre-school (3–6 years)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
School Age (6–12 years)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Adolescent (12–18 years)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Young Adult (18–30 years)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Mature Adult (30–60 years)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4
Elderly (>60 years)	☐ 1	☐ 2	☐ 3	☐ 4	☐ 1	☐ 2	☐ 3	☐ 4

Work History

Years Experience in Clinical Specialty:

Most recent facility worked at:

☐ I hereby certify that ALL information I have provided to Clover Health Services, on this skills checklist and all other documentation, is true and accurate. I understand and acknowledge that any misrepresentation or omission may result in disqualification from employment and/or immediate termination.